

We claim.

1. An inhibitor of catalytically active memapsin 2 which binds to the active site of the memapsin 2 defined by the presence of two catalytic aspartic residues and substrate binding cleft.

5 2. The inhibitor of claim 1 comprising an isostere of the active site of memapsin 2.

3. The inhibitor of claim 2 comprising a molecule having the general form $X-L_4-P_4-L_3-P_3-L_2-P_2-L_1-P_1-L_0-P_1'-L_1'-P_2'-L_2'-P_3'-L_3'-P_4'-L_4'-Y$, wherein P_x represent the substrate specificity position relative to the
10 cleavage site which is represented by an $-L_0-$, and L_x represent the linking regions between each substrate specificity position, P_x , and
wherein L_0 is a non-hydrolyzable bond and P_1' is $-R_1CR_3-$, wherein R_1 is a group smaller than CH_2OH (side chain of serine), and at least two other P positions are a hydrophobic group.

15 4. The inhibitor of claim 3 which is OM99-1.

5. The inhibitor of claim 3 which is OM99-2.

6. The inhibitor of claim 3 having the structure of Figure 11.

7. The inhibitor of claim 3 having the structure of Figure 12.

8. The inhibitor of claim 3 having the structure of Figure 13.

20 9. The inhibitor of claim 3 having the structure of Figure 14.

10. The inhibitor of claim 1 having an K_i of less than or equal to 10^{-7} M.

11. The inhibitor of claim 1 which binds to crystallized enzyme characterized by the parameters in Table 2 when bound to OM-99-2.

25 12. The inhibitor of claim 11 having a K_i of less than or equal to 10^{-6} M.

13. The inhibitor of claim 11 having a K_i of less than or equal to 2 nM.

30 14. The inhibitor of claim 13 having a K_i of less than or equal to 1 nM.

15. The inhibitor of claim 11 having a root mean square difference of less than or equal to 0.5 Å for the side chain and backbone atoms for amino acids 18-379 of memapsin 2.

16. The inhibitor of claim 1 which is permeable to the blood brain
5 barrier.

17. The inhibitor of claim 1 which blocks cleavage by memapsin 2 under physiological conditions.

18. The inhibitor of claim 1 which is a non-amino acid small molecule.

19. The inhibitor of claim 18 having a molecular weight of less than
10 800 Daltons.

20. A method of synthesis of a Leu*Ala dipeptide isostere.

21. A method for treating a patient to decrease the likelihood of
15 developing or the progression of Alzheimer's disease comprising administering to the individual an effective amount of an inhibitor of memapsin 2 having an K_i of less than or equal to 10^{-7} M or which binds to crystallized enzyme characterized by the parameters in Table 2 when bound to OM-99-2.

22. The method of claim 21 wherein the inhibitor is administered orally.

23. The method of claim 21 wherein the inhibitor blocks cleavage of
20 APP.

ADD
C1

add D6
add E1